Copper Oxychloride	Group	M01	Fungicide	j
Copper Hydroxide	Group	M01	<u>F</u> ungicide	j

FIFRA SECTION 24(C) SPECIAL LOCAL NEED REGISTRATION FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS







EPA Reg. No. 10163-402 EPA SLN No. TX-230003

WARNING / AVISO

This label for Badge X2 expires and must not be distributed or used in accordance with this SLN registration after 12/31/2028.

DIRECTIONS FOR USE

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- Labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

CROP	DISEASE	APP. RATE (LBS PRODUCT/A)	MAX. APP. RATE/YEAR (LBS PRODUCT/A)	COMMENTS
Organic Rice	Brown Spot (Cochliobolus miyabeanus) and Narrow Brown Leaf Spot (Cercospora janseana)	1.15 – 1.8	3.6	 Apply 1.15 – 1.8 lbs/A of Badge X2 either once at boot stage and heading, or once at boot stage only. For susceptible cultivars or when environmental conditions are favorable for disease on emerging panicles, use higher rates. Ensure that the spray is applied uniformly to achieve effectiveness. To minimize risk of crop injury, apply to dry foliage. Do not apply this product in a spray solution with a pH level below 7.0 to prevent crop injury.
	GROUND APPLICATION: mix 1.15 – 1.8 lbs. of product (maximum of 1.8 lbs.) in a minimum of 10 – 20 gal. of water per acre. AERIAL APPLICATION: mix 1.15 – 1.8 lbs. of product (maximum of 1.8 lbs.) in a minimum of 5 - 10 gal. of water per acre. PRECAUTIONS: Badge X2 is most effective when applied and allowed to dry before any rainfall. For optimal results, make two applications of Badge X2 during the growing season. Apply to dry foliage to minimize the risk of crop injury.			

RESTRICTIONS:

- Do not exceed 3.6 lbs of Badge X2 per acre per year.
- Do not exceed 1.8 lbs of Badge X2 per acre per application.
- Do not make more than two applications of Badge X2 during the growing season.
- Do not apply this product in a spray solution with a pH level below 7.0 to prevent crop injury.

Section 24(c) Registrant: Gowan Company P.O. Box 5569 Yuma, AZ 85366-5569

Rev 04/06/2023