

Supplemental Labeling

Sandea[®]
Herbicide

Sandea[®] is a selective herbicide for control of listed weeds including nutsedge

EPA REG. No. 81880-18-10163

ACTIVE INGREDIENT:	% BY WT.
Halosulfuron-methyl, methyl 3-chloro-5-(4,6-dimethoxypyrimidin-2-ylcarbamoylsulfamoyl)-1-methylpyrazole-4-carboxylate.....	75.0%
OTHER INGREDIENTS	25.0%
	TOTAL 100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Read the entire label before using this product.

- This labeling must be in the possession of the user at the time of pesticide application.
- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

AERIAL APPLICATIONS IN SUGARCANE – FLORIDA ONLY

Aerial Applications:

Apply this product or approved tank mixtures with properly calibrated equipment in 3-15 gallons of water per acre. Thoroughly clean equipment prior to mixing spray solution. Avoid streaking, skips, overlaps, and spray drift during applications.

SUGARCANE (30)	2/3 - 1 1/3	<p>When used alone, apply SANDEA prior to planting, prior to emergence, after the emergence of the sugarcane, and until row closure. Mechanical cultivation may be required to control weed species not on the label. If so, a sequential treatment may be required to control weeds in areas of disturbed soil.</p> <p>Apply SANDEA at 2/3 -1 1/3 ounces by weight per acre (0.031-0.062 pound active ingredient per acre) in combination with glyphosate agricultural herbicides for pre-plant burn down of emerged annual grasses, broadleaf weeds, and nutsedge in sugarcane.</p> <p style="text-align: center;">Tank Mixtures for Sugarcane</p> <p>Tank mixtures with SANDEA can include, but are not limited to Asulox[®], Atrazine, Callisto[®], Envoke[®], Evik[®], glyphosate, or 2,4-D.</p> <p>Refer to the specific product labels and observe all precautions, mixing and application instructions, and follow-crop intervals for all products used in tank mixtures.</p> <ul style="list-style-type: none"> • Refer to the "ROTATIONAL CROP INFORMATION" section of this label for applicable rotational crop restrictions. • No more than 3 applications (including pre-plant applications) may be made with the total use rate not to exceed 2 2/3 ounces of product by weight (0.125 pound active ingredient) per acre per year. • Following application to foliage allow 30 days before grazing domestic livestock, harvesting forage, or harvesting silage. • Consult "Use Precautions" and "For Optimum Results" sections for important usage information.
-----------------------	-------------	---