


GROUP	21	ACARICIDE
-------	----	-----------

**FIFRA §2(ee) RECOMMENDATION
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF FLORIDA, GEORGIA, MAINE,
MICHIGAN, NEW JERSEY AND NORTH CAROLINA.**



EPA REG. NO. 10163-322

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.
- This recommendation is made as permitted under FIFRA §2(ee) and has not been submitted to or approved by the EPA.

CROP	PEST	RATE FL OZ/ACRE	RATE LB A.I./ACRE	COMMENTS
BUSHBERRY SUBGROUP 13-07B (7)  Aronia berry; blueberry, highbush; blueberry, lowbush; buffalo currant; Chilean guava; cranberry, highbush; currant, black; currant, red; elderberry; European barberry; gooseberry; honeysuckle, edible; huckleberry; jostaberry; Juneberry (Saskatoon berry); lingonberry; native currant; salal; sea buckthorn; cultivars, varieties, and/or hybrids of these	Blueberry bud Mite, Southern Red Mite	24 -36	0.3 – 0.48	Apply in at least 50 gallons of water per acre. Use higher rates for heavier mite pressure.
				<ul style="list-style-type: none"> • Do not make more than one application per year. • Do not exceed application rate of 36 fl oz of product (0.48 lb a.i.) per acre per year. • Do not apply this product until crop is through petal fall unless the conditions under the Direction For Use section on the Section 3 label have been met.

RESTRICTIONS ON ALL CROPS:

- Aerial applications are prohibited.
- Do not apply by any type of irrigation system.
- Only ground sprayer and airblast applications are permitted.
- Do not apply during conditions when product will drift to other crops and non-target areas.
- Do not plant rotational crops, other than those on this label, within 30 days of this application. Do not plant root, tuber and bulb vegetables within 120 days of this application.