

FIFRA §2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF FLORIDA and TEXAS

GROUP 21A ACARICIDE



EPA REG. NO. 81880-28-10163

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.
- This recommendation is made as permitted under FIFRA §2(ee) and has not been submitted to or approved by the EPA.

CROP	PEST	RATE FL OZ/ACRE	COMMENTS
Citrus fruit group 10-10 (7 day PHI)	Diaprepes root weevil (<i>Diaprepes abbreviatus</i>) Apopka weevil (<i>Diaprepes abbreviatus</i>)	16 - 17	Apply 16-17 oz in sufficient water to achieve thorough coverage. Use the higher rate of to ensure adequate concentration in full size trees with dense foliage. When combining with summer oils, use a minimum of 5 gallons of oil and 11 oz of NEXTER SC per acre. In Florida Only - may be applied in low volume application equipment with a minimum water volume of 20 gallons of water per acre. It is the user's responsibility to ensure thorough spray coverage in these low volume applications.
<ul style="list-style-type: none"> • Do not make more than two applications per year. • Do not apply more than 17 oz per acre per application • Allow a minimum of 30 days between sequential applications. • For rates above 16 oz per acre, apply on a 90 day interval. 			