

FIFRA SECTION §24(C) SPECIAL LOCAL NEED LABEL

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

THIS LABEL IS VALID UNTIL DECEMBER 31, 2024 UNLESS OTHERWISE AMENDED, WITHDRAWN, CANCELED, OR SUSPENDED

EPTAM® 7E

Selective Herbicide - Emulsifiable Liquid

EPA Reg. No. 10163-283, EPA SLN No. OR-190006

FOR CONTROL OF WEEDS IN GRASS GROWN FOR SEED – ESTABLISHED STANDS.

ACTIVE INGREDIENT:

EPTC: S-ethyl dipropylthiocarbamate.....87.8%

INERT INGREDIENTS:.....12.2%

Total 100.0%

Contains 7 pounds active ingredient per gallon.
Contains petroleum distillates

**KEEP OUT OF REACH OF CHILDREN
WARNING-AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

ENVIRONMENTAL HAZARDS

This chemical is toxic to mammals. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- Follow all applicable directions, restrictions, and precautions, including statements pertaining to the Worker Protection Standards, on the EPA-registered main label and this label.
- This label must be in the possession of the user at the time of pesticide application.

CROP	RATE Pints/Acre	COMMENTS
GRASS GROWN FOR SEED – ESTABLISHED STANDS (such as perennial ryegrass, orchardgrass, tall fescue, fine fescue, bluegrass and bentgrass)	3.5 - 5	PACIFIC NORTHWEST: Apply broadcast and incorporate into the top ½ - 1 inch of dry soil via rainfall or irrigation in the amount of 0.25 to 0.5 inch. Rain events greater than 1 inch may produce undesirable control and crop injury or stunting. Timing: Fall: Apply Eptam in a broadcast spray at a timing prior to weed seed germination and within 36 hours of a predicted rain event when grass seed crops are actively growing. In perennial ryegrass only, fall applications can be made to newly established stands once the 1 st tiller of the crop has developed. Eptam can be applied to spring planted tall fescue during the fall in the establishment year. All other grass seed crops need to have one seed crop harvested before applying Eptam 7E. Do not apply after December 1 st . OR Spring: Apply Eptam in a broadcast spray at a timing prior to weed seed germination and within 7 days of a predicted rain event when grass seed crops are actively growing. Applications should be made when established grass seed crops have 4-6 tillers. Tank mix with a postemergence herbicide to control germinated weeds.
		<ul style="list-style-type: none"> • Do not enter or allow worker entry during the restricted-entry interval of 2 days. • Do not make more than one application per year. • Eptam 7E does not control germinated or established annual weeds present at application. • Eptam 7E is recommended for use on mineral soils only (soils containing less than 10% organic matter). • Application must be made to a dry soil surface (at least ½ inch deep) free from dew and incidental moisture. • For optimum weed control, incorporation should be done as soon as possible and no later than 36 hours following application. Postharvest residue, including straw and stubble, should be removed from the field prior to application.

24(c) Registrant:
Gowan Company
P.O. Box 5569
Yuma, AZ 85366