

FIFRA §2(ee) RECOMMENDATION FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF CALIFORNIA AND ARIZONA

BENSULIDE	GROUP	0	HERBICIDE
-----------	-------	---	-----------

Prefar® 4-E

Selective Herbicide

EPA REG. NO. 10163-200

Herbicide for Control of Additional Pest: Corn Spurry in Listed Crops

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.
- This recommendation is made as permitted under FIFRA §2(ee) and has not been submitted to or approved by the EPA.

PESTS CONTROLLED: Corn Spurry

CROP	BROADCAST RATE Quarts/Acre	COMMENTS
CUCURBIT VEGETABLE GROUP , including but not limited to: Chinese Waxgourd (Chinese Preserving Melon), Cucumbers, Gherkin, Edible Gourd (Chinese Okra), Muskmelons (including Cantaloupe, Honeydew Melon), Summer Squash, Winter Squash, Watermelons	5 - 6	Apply preplant or preemergence to flat-planted or bedded crops. Follow application instructions given on this label.
BRASSICA (COLE) LEAFY VEGETABLE GROUP , including but not limited to: Broccoli, Broccolini, Broccoflower, Chinese Broccoli, Broccoli Raab (Rapini), Brussels Sprouts, Cabbage, Chinese Cabbage (Bok Choy, Napa), Chinese Mustard Cabbage (Gai Choy), Cauliflower, Cavalo Broccolo, Collards, Kale, Kohlrabi, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens	5 - 6	Do not apply more than 6 lbs. a.i. per acre per season. Apply preplant or preemergence. Follow application instructions given on this label.
FRUITING VEGETABLES Eggplant, Pepinos, Peppers (Bell Peppers, Chili Peppers, Cooking Peppers, Pimentos, Sweet Peppers), Tomatillo	5 - 6	Apply preplant or preemergence. Follow application instructions given on this label.

<p>LEAFY VEGETABLES (EXCEPT BRASSICA VEGETABLE) GROUP, including but not limited to: Arugula (Roquette), Cardoon, Celery, Chinese Celery, Chervil, Cress (Garden, Upland), Dock (Sorrel), Endive (Escarole), Florence Fennel, Lettuce (Head or Leaf), Parsley, Radicchio (Red Chicory)</p>	<p>5 - 6</p>	<p>Do not apply more than 6 lbs. a.i. per acre per season. Apply preplant or preemergence. Follow application instructions given on this label. Do not use Prefar 4-E on spinach or Swiss Chard as severe phytotoxicity will occur.</p>
<p>DRY BULB VEGETABLES: Garlic, Bulb Onions, Shallots (Do not use in the Willamette Valley of Oregon)</p>	<p>5 - 6</p>	<p>Do not apply more than 6 lbs. a.i. per acre per season. Apply preplant or preemergence. Follow application instructions given on this label. Onions germinate and grow best when planted in fine-textured soil with good water drainage where the soil salinity value is less than 4 ECe* and soil temperature is between 55° and 75°. Use of Prefar on onions under less favorable soil and environmental conditions may reduce stand and vigor resulting in lower yield. To avoid injury, do not incorporate by mechanical methods in California or Arizona. Consult your local extension agent or university specialist for further information. (*ECe is the electrical conductivity extract.)</p>

Prefar® is a registered trademark of Gowan Company, L.L.C.

12-8-17