

**FIFRA §2(ee) RECOMMENDATION
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF DELAWARE,
FLORIDA, GEORGIA, MARYLAND, MICHIGAN, NEW JERSEY, NORTH
CAROLINA, OHIO, PENNSYLVANIA, SOUTH CAROLINA, VIRGINIA**

QUINTEC[®] FUNGICIDE

For suppression of Bacterial Spot in Peppers

- This labeling must be in the possession of the user at the time of pesticide application.
- It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

Quintec aids in the suppression of bacterial spot in peppers when used in an alternation or tank mix spray program with other products that have activity on bacterial spot.

Use 6 oz of product per acre in no less than 30 gallons of water per acre.

Refer to the label affixed to the container for Quintec for general use precautions and restrictions, mixing instructions, and application guidelines. Refer to the supplemental label for Quintec for use on peppers for rotational crop restrictions, disease resistance management, application methods, and specific use restrictions. Read and follow all applicable directions for use, precautions and limitations.

Resistance Management for *Xanthomonas*

Quintec must be alternated with other product recommended for effective control of *Xanthomonas*. When more than one application of Quintec is made per crop, at least one of the applications of Quintec must be a tank mixture with a product effective on *Xanthomonas*. Do not make more than two consecutive applications of Quintec before alternating to another product which is effective on *Xanthomonas*. Consult your local or state agriculture authorities for resistance management strategies that are appropriate for your disease management program.

Resistance Management of Powdery Mildew

If powdery mildew is present, the number of sprays of Quintec per crop must not exceed 50% of the total number of powdery mildew sprays. Quintec is a Group 13 fungicide. Alternation with effective fungicides from other fungicide Groups is recommended after each application of Quintec. When more than one application is made per crop, at least one of the applications of Quintec must be a tank mixture with a product that is effective on powdery mildew and has a different fungicide Group number than Quintec. Apply the alternate product or tank mix within 14 days of applying Quintec. Do not make more than two consecutive applications of Quintec.

Quintec[®] is a registered trademark of Nissan Chemical Corporation

EPA Reg. No. 33906-26-10163

20-10-03



Distributed by
Gowan Company
P. O. Box 5569
Yuma, AZ 85366-5569