

**FIFRA §2(ee) RECOMMENDATION  
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF DELAWARE, MARYLAND,  
MASSACHUSETTS, NEW JERSEY, PENNSYLVANIA, VIRGINIA AND WEST VIRGINIA**

**SPOTTED LANTERNFLY IN GRAPES, PEACHES AND NECTARINES**



**EPA REG. NO. 10163-317**

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered labels are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

**DIRECTIONS FOR USE**

<b>CROP</b>	<b>PEST</b>	<b>RATES</b>	<b>COMMENTS</b>
Grapes	Spotted Lanternfly (SLF)	<b>Foliar:</b> 5 fl oz/A 0.13 lbs ai/A	Higher water volumes provide improved insect control.  Begin application when pest activity is first noticed or when insects reach threshold levels per State and County Extension Service recommendations. Repeat as needed to maintain control, but not more often than every 7 days. For best results, time application before a damaging population becomes established.
	Spotted Lanternfly (SLF)	<b>Soil:</b> 10.5 fl oz/A 0.27 lbs ai/A	SCORPION may be mixed and/or alternated with commonly used insecticides to comply with local IPM and resistance management programs.
<p><b>Restriction: Regardless of application method, do not apply more than a total of 20.9 fl oz/A of SCORPION (0.529 lb ai/A) per season.</b></p> <p><b>Foliar Application</b></p> <ul style="list-style-type: none"> <li>• Apply with air or ground equipment in adequate water for uniform coverage (Do not use less than 5 gallons/acre for aerial applications or 10 gallons/acre for ground applications).</li> <li>• Do not apply SCORPION within one (1) day of harvest.</li> <li>• Do not apply more than a total of 10.25 fl oz/A of SCORPION (0.259 lb ai/A) per season.</li> </ul> <p><b>Soil Application</b></p> <ul style="list-style-type: none"> <li>• Make only one (1) soil application per season.</li> <li>• Apply with ground equipment in adequate water for uniform coverage (10 to 100 gals/A).</li> <li>• Do not apply SCORPION within twenty-eight (28) days of harvest.</li> <li>• Do not apply more than a total of 10.5 fl oz/A of SCORPION (0.266 lb ai/A) per season.</li> <li>• For drip application, prior to injection, mix specified dosage in sufficient carrier volume (minimum of 2 gals of water per 1 lb of product) to ensure uniform application and incorporation into the soil using drip or trickle irrigation water. Apply towards the end of the irrigation run to ensure the product does not leach past the root zone.</li> </ul>			



The Go To Company

P.O. Box 5569 ▲ Yuma, AZ 85366-5569 ▲ Phone (928) 783-8844 ▲ FAX (928) 343-9255

CROP	PEST	RATE	COMMENTS
Peach Nectarine	Spotted Lanternfly (SLF)	<b>Foliar:</b> 5.25 to 7 fl oz/A (0.135 to 0.180 lb ai/A)	<p>Higher water volumes provide improved insect control.</p> <p>Begin applications when first pest activity is noticed or when insects reach threshold levels per State and County Extension Service recommendations. Repeat as needed to maintain control, but not more often than every 7 days. For best results, time application before a damaging population becomes established.</p> <p>Under severe pest pressure, use the higher specified rates.</p> <p>The rate applied affects the length of control. Use the high rate where infestations occur later in crop development, or where pest pressure is continuous.</p> <p>SCORPION 35SL INSECTICIDE may be mixed and/or alternated with commonly used insecticides to comply with local IPM and resistance management programs.</p>
	Spotted Lanternfly (SLF)	<b>Soil:</b> 10.5 fl oz/A (0.270 lb ai/A)	
<p><b>NOTE: Regardless of application method do not apply more than a total of 14.25 fl oz/A of SCORPION (0.360 lb. ai) per acre per season.</b></p> <p><b>Foliar Application</b></p> <ul style="list-style-type: none"> <li>• Apply with air or ground equipment in adequate water for uniform coverage (A minimum of 5 gals/A by air or 50 gals/A by ground).</li> <li>• Do not apply SCORPION within three (3) days of harvest.</li> <li>• Do not apply more than a total of 10.5 fl oz/A of SCORPION (0.270 lb ai) per acre per season.</li> <li>• Interval between applications cannot be less than 7 days.</li> </ul> <p><b>Soil Application</b></p> <ul style="list-style-type: none"> <li>• Do not apply SCORPION within twenty one (21) days of harvest</li> <li>• Apply with ground equipment in adequate water for uniform coverage (A minimum of 100 gals/A).</li> <li>• Do not apply more than a total of 10.5 fl oz/A of SCORPION (0.270 lb ai) per acre per year.</li> </ul> <p>Apply specified dosage in sufficient carrier volume to insure uniform application and distribution within and around the root zone of each tree using one of the following methods:</p> <ol style="list-style-type: none"> <li>1. As a drench. Applications should be made with sufficient water to insure incorporation into the root zone.</li> <li>2. Using drip, trickle, micro sprinkler or any customized irrigation system derived from those systems to water trees independently.</li> </ol>			

Scorpion® is a registered trademark of Gowan Company, L.L.C.

18-08-01