

**FIFRA §2(ee) RECOMMENDATION
FOR DISTRIBUTION AND USE ONLY WITHIN THE STATES OF
WASHINGTON, OREGON**



For Agricultural Use Only

EPA REG. NO. 8033-103-10163

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions, and precautions on the EPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.
- This recommendation is made as permitted under FIFRA §2(ee) and has not been submitted to or approved by the EPA.

POME FRUIT (within Crop Group 11)

SITE	DISEASE	TORINO FUNGICIDE (oz/acre) [POUNDS ACTIVE]	USE DIRECTIONS	RESTRICTIONS
POME FRUIT apple, pear, crabapple, loquat, mayhaw, Quince, Oriental pear	Powdery mildew	1 application at 5.5-6.8 [0.036-0.044]	Begin application at first sign of disease development. For the control of other foliar diseases of pome fruit applications of registered fungicides must be made according to their label directions for use.	The minimum retreatment interval is fourteen (14) days. Do not make more than one (1) application at the 6.8 oz. product rate per calendar year. Do not exceed a total of 6.8 oz of product (0.044 lbs. active ingredient) per acre per calendar year. Do not apply within 14 days of harvest; (PHI = 14 days).